

MEDICATION GUIDE

DURAGESIC[®] (Dur-ah-GEE-zik)

(FENTANYL
TRANSDERMAL
SYSTEM) 

IMPORTANT:

- Keep DURAGESIC[®] in a safe place away from children and pets. Accidental use by a child or pet is a medical emergency and may result in death. If a child or pet accidentally uses DURAGESIC[®], get emergency help right away.
- Make sure you read the separate **“Instructions for Applying a DURAGESIC[®] Patch.”** Always use a DURAGESIC[®] Patch the right way. DURAGESIC[®] can cause serious breathing problems and death, especially if it is used the wrong way.
- DURAGESIC[®] is a federally controlled substance (C-II) because it can be abused. Keep DURAGESIC[®] in a safe place to prevent theft. Selling or giving away DURAGESIC[®] may harm others, and is against the law.
- Tell your doctor if you (or a family member) have ever abused or been dependent on alcohol, prescription medicines or street drugs.

Read the Medication Guide that comes with DURAGESIC[®] before you start using it and each time you get a new prescription. There may be new information. This Medication Guide does not take the place of talking to your healthcare provider about your medical condition or your treatment. Make sure you read and understand all the instructions for using DURAGESIC[®]. Do not use DURAGESIC[®] unless you understand everything. Talk to your healthcare provider if you have questions.

What is the most important information I should know about DURAGESIC[®]?

DURAGESIC[®] is a skin patch that contains fentanyl. Fentanyl is a very strong opioid narcotic pain medicine that can cause **serious and life-threatening breathing problems**. Serious and life-threatening breathing problems can happen because of an overdose or if the dose you are using is too high for you. Call your doctor right away or get emergency medical help if you:

- have trouble breathing, or have slow or shallow breathing
- have a slow heartbeat
- have severe sleepiness
- have cold, clammy skin
- feel faint, dizzy, confused, or cannot think, walk, or talk normally
- have a seizure
- have hallucinations

DURAGESIC[®] is only for adults and children over the age of two with persistent, moderate to severe chronic pain and who:

- are already using another strong opioid narcotic pain medicine around-the-clock, and have been using the medicine regularly for a week or longer. This is called being opioid-tolerant.

- have pain that cannot be controlled with other medicines

Do not use DURAGESIC[®]:

- if you are not already using another opioid narcotic medicine and are not opioid tolerant
- if you need opioid pain medicines for only a short time
- for pain from surgery, medical or dental procedures
- if your pain can be taken care of by occasional use of other pain medicines
- in children who are less than 2 years of age
- if you have asthma symptoms or have severe asthma

A DURAGESIC[®] patch must be used only on the skin of the person for whom it was prescribed. If the patch comes off and accidentally sticks to the skin of another person, take the patch off of that person right away, wash the area with water, and get medical care for them right away.

DURAGESIC[®] patch is not safe for everyone. Tell your doctor about all of your medical conditions.

Tell your doctor if you are planning to become pregnant, are pregnant, or

breastfeeding. DURAGESIC® may cause serious harm to a baby.

Tell your doctor about all the medicines you take. Some medicines may cause serious or life-threatening side effects when used with DURAGESIC®. Your doctor will tell you if it is safe to take other medicines while you are using DURAGESIC®.

Know the medicines you take. Keep a list of your medicines to show to your doctor and pharmacist.

How should I use DURAGESIC®?

Read the separate “Instructions for Applying a DURAGESIC® Patch”.

• **You must always use DURAGESIC® patches the right way:**

- **Do not** use a DURAGESIC® patch if the pouch seal is broken, or the patch is cut, damaged, or changed in any way.
- **Do not** use heat sources such as heating pads, electric blankets, heat lamps, tanning lamps, saunas, hot tubs, or heated waterbeds while wearing a DURAGESIC® patch.
- **Do not** take hot baths or sunbathe while wearing a DURAGESIC® patch.

• **If you have problems with the DURAGESIC® patch not sticking:**

1. Apply first aid tape only to the edges of the patch.
2. If problems with the patch not sticking persist, cover the patch with Bioclusive™ or Tegaderm™. These are special see-through adhesive dressings. **Never cover a DURAGESIC® patch with any other bandage or tape.**

• **If your DURAGESIC® patch falls off before 3 days or 72 hours, fold the sticky side together and flush down a toilet. Put a new one on at a different skin site.**

• **Do not change your dose unless your doctor tells you to.** Your doctor may change your dose after seeing how the medicine affects you. Do not use DURAGESIC® more often than prescribed. Call your doctor if your pain is not well controlled while using DURAGESIC®.

• **Do not stop using DURAGESIC® suddenly.** Stopping DURAGESIC® suddenly can make you sick with withdrawal symptoms (for example, nausea, vomiting, diarrhea, anxiety, and shivering). Your body can develop a physical dependence on DURAGESIC®. If

your doctor decides you no longer need DURAGESIC®, ask how to slowly reduce this medicine so you don't have withdrawal symptoms. Do not stop taking DURAGESIC® without talking to your doctor.

• **Do not wear more than one DURAGESIC® patch at a time,** unless your doctor tells you to do so.

• **Call your doctor right away if**

• **You get a fever higher than 102°F**

• Your body temperature increases from exercise

A fever or increase in body temperature may cause too much of the medicine in DURAGESIC® to pass into your body.

• **If you use more DURAGESIC® than your doctor has prescribed, get emergency medical help right away.**

• **Do not drink any alcohol while using DURAGESIC®.** Alcohol can increase your chances of having serious side effects.

• **Do not drive, operate heavy machinery, or do other possibly dangerous activities** until you know how DURAGESIC® affects you. DURAGESIC® can make you sleepy. Ask your doctor to tell you when it is okay to do these activities.

• When you remove your DURAGESIC® patch, fold the sticky sides of a used DURAGESIC® patch together and flush it down the toilet. **Do not put used DURAGESIC® patches in a trash can.**

What are the possible side effects of DURAGESIC®?

Serious side effects include:

• **Life-threatening breathing problems.** See “What is the most important information I should know about DURAGESIC®?”

• **Low blood pressure.** This can make you feel dizzy if you get up too fast from sitting or lying down.

The common side effects with DURAGESIC® are nausea, vomiting, constipation, dry mouth, sleepiness, confusion, weakness, sweating, and pain and redness where the patch was applied.

Constipation is a very common side effect of all opioid medicines. Talk to your doctor about the use of laxatives and stool softeners to prevent or treat constipation while taking DURAGESIC®.

Talk to your doctor about any side effect that

concerns you.

These are not all the possible side effects of DURAGESIC[®]. For a complete list, ask your doctor or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store DURAGESIC[®]?

- Store in original unopened pouch at room temperature.
- Keep a DURAGESIC[®] patch in its protective pouch until you are ready to use it.
- **Keep DURAGESIC[®] in a safe place out of the reach of children and pets.**
- Dispose of DURAGESIC[®] patches you no longer need. Open the unused packages, fold the sticky sides of the patches together, and flush them down the toilet.

General information about the safe and effective use of DURAGESIC[®]

- Do not use DURAGESIC[®] for a condition for which it was not prescribed.
- **Do not give DURAGESIC[®] to other people, even if they have the same symptoms you have. DURAGESIC[®] can harm other people and even cause death. Sharing DURAGESIC[®] is against the law.**

This Medication Guide summarizes the most important information about DURAGESIC[®]. If you would like more information, talk with your

doctor. You can ask your doctor or pharmacist for information about DURAGESIC[®] that is written for doctors.

For questions about DURAGESIC[®], call the Ortho-McNeil Janssen Scientific Affairs Customer Communications Center at 1-800-526-7736. If this is a medical emergency, please call 911.

What are the ingredients of DURAGESIC[®]?

Active Ingredient: fentanyl

Inactive ingredients: polyester/ethyl vinyl acetate film backing, polyacrylate adhesive.

This Medication Guide has been approved by the United States Food and Drug Administration.

Bioclusive[™] is a trademark of Ethicon, Inc.
Tegaderm[™] is a trademark of 3M

Manufactured by:
ALZA Corporation
Vacaville, CA 95688

Manufactured for:
PriCara[®], Division of Ortho-McNeil-Janssen
Pharmaceuticals, Inc.
Raritan, NJ 08869

Insert New Code

July 2009

© Ortho-McNeil-Janssen Pharmaceuticals, Inc.
2009